

Resonances in the current-voltage characteristics of a dissipative Josephson junction

J. Müllers and A. Schmid

*Institut für Theorie der Kondensierten Materie, Universität Karlsruhe, 76128 Karlsruhe,
Germany*

(February 1, 2008)

Abstract

The I-V characteristics of a Josephson junction shunted by an Ohmic resistor shows sharp peaks when levels in neighbouring wells are crossing. We consider the shape and size of these peaks using a double-well model where the wells are given in parabolic approximation. The friction of arbitrary strength is included by the help of a bath of infinitely many degrees of freedom.

74.50.+z

Keywords: Josephson junction; Caldeira-Leggett model; dissipation; tunneling

I. INTRODUCTION

The past decade has seen a considerable interest and remarkable activity in an area which presently is often referred to as macroscopic quantum mechanics. Specifically, one has been interested in quantum phenomena of macroscopic objects [1].

In particular, macroscopic quantum tunneling [2] (quantum decay of a metastable state), and quantum coherence [3] have been studied. Soon, it became clear that dissipation has a profound influence on these quantum phenomena. Phenomenologically, dissipation is the consequence of an interaction of the object with an environment which can be thought of as consisting of infinitely many degrees of freedom. Specifically, the environmental degrees of freedom may be chosen to be harmonic oscillators such that we may consider the dissipation as a process where excitations, that are phonons, are emitted and absorbed. Such a model (Caldeira-Leggett model) has been the basis in Ref. [2] where the influence of dissipation on tunneling has been explored.

As far as quantum coherence is concerned, the most simple system is an object with two different quantum states: it is thought to represent the limiting case of an object in a double-well potential where only the lowest energy states in each of the two wells is relevant and where the tunneling through the separating barrier allows for transitions that probe the coherence. Since a two-state system is equivalent to a spin-one-half problem, this standard system is often referred to by this name. In particular, with the standard coupling to a dissipative environment made of harmonic oscillators, it is called the spin-boson problem which has been studied repeatedly in the past [3,4].

Level quantization and resonant tunneling have been observed recently [5] in a double-well quantum-dot system. However, the influence of dissipation was not considered in this experiment. On the other hand, it seems that Josephson junctions are also suitable systems for obtaining experimental evidence pertaining to macroscopic quantum effects. In this

context, evidence for level quantization [6], [7] and for quantum decay [8] have been obtained; however, macroscopic quantum coherence has evaded observation so far.

Specifically, a Josephson junction may be characterized by a current-phase relation

$$I(\varphi) = I_J \sin \varphi , \quad (1)$$

where the phase φ is related to the voltage difference U by

$$\hbar \dot{\varphi} = 2eU . \quad (2)$$

Therefore, the phase of a Josephson junction shunted by a capacitance C and biased by an external current I_x obeys a classical type of equation of motion

$$M \ddot{\varphi} = - \frac{\partial V(\varphi)}{\partial \varphi} , \quad (3)$$

with the mass

$$M = \left(\frac{\hbar}{2e} \right)^2 C \quad (4)$$

and the potential energy

$$V(\varphi) = - \frac{\hbar}{2e} [I_J \cos \varphi + I_x \varphi] . \quad (5)$$

A widely discussed model of a dissipative object is the one where the Josephson junction is also shunted by an Ohmic resistor R . In this case, the classical equation of motion (3) has to be replaced by

$$\begin{aligned} M \ddot{\varphi} &= - \frac{\partial V(\varphi)}{\partial \varphi} - \eta \dot{\varphi} , \\ \eta &= \left(\frac{\hbar}{2e} \right)^2 \frac{1}{R} . \end{aligned} \quad (6)$$

The model of a dissipative environment according to the above specification has been discussed by Caldeira and Leggett [2].

The potential energy $V(\varphi)$ of Eq. (5) displays wells at $\varphi \simeq 2n\pi$ with depth shifted by an amount $\Delta \simeq (2\pi\hbar/2e)I_x$. If the wells are sufficiently deep, one needs to concentrate only

on transitions between pairs of adjacent wells. Thus, one arrives at the double well problem mentioned above.

The analysis in this paper goes beyond the limiting situation where only the lowest level in each of the two wells is of importance. Roughly, this is realized when the level separation $\hbar(2E_J/M)^{1/2} \simeq (2e\hbar I_J/C)^{1/2}$ is smaller than or comparable with Δ . In particular, we will concentrate on resonance phenomena which are expected to show up whenever two levels in the adjacent wells happen to cross when the bias current I_x , that is Δ , is varied.

For such values of the bias current, there appear sharp asymmetric peaks in the current-voltage characteristic of the Josephson junction. This phenomenon has been studied by one of us (A.S., together with A.I. Larkin and Yu. N. Ovchinnikov) [9] within the standard model in the one-phonon approximation. For bias currents that correspond to crossings of the next and next nearest levels (e.g. ground state in the left well and the first or second excited state at the right side), it is possible to neglect processes in the reverse direction provided that the temperature is sufficiently low. Thus, the restriction to a double well system receives additional support.

The transfer of the object from the left to the right potential well is accompanied by the emission of an infinite number of phonons. Therefore, in a recent paper by Ovchinnikov and one of us (A.S.) [10] the fact is taken into account that in the resonance region, the contribution of phonons of small energy is important as well as the contribution of resonance phonons with energy equal to the distance between levels in the wells.

The above analysis relies on a perturbation theory for weak dissipation. Moreover, the calculations in that article are rather intricate. Therefore, we want to reconsider the problem in a more transparent way and thereby get rid of the restriction of small dissipation.

II. DESCRIPTION OF THE MODEL HAMILTONIAN AND ITS EIGENSTATES

The model consists of a particle, called "object" (coordinate R_1), which is coupled (in the sense of Caldeira and Leggett [2]) to a "bath" of harmonic oscillators (coordinates R_j).

We shall use the conventions $j \in \{2, \dots, N\}$ for the bath oscillators and $k \in \{1, \dots, N\}$ for the indices of all coordinates in the model. The double-well potential is approximated by two parabolas about the minima of the two wells as it is depicted in fig. 1.

The phase φ of the Josephson contact then corresponds to the object coordinate R_1 of the model, and the voltage U is related to the tunneling rate J by $2eU = \dot{\varphi} = 2\pi J$. As it has already been remarked, the current I_x is proportional to the bias Δ of the two wells. Thus, calculating the transition rate for different values of the bias Δ is equivalent to the determination of the I-V characteristics.

Specifically, we want to write the Hamiltonian of the model in the form

$$\begin{aligned}\hat{H} &= \frac{1}{2m} \sum_k \hat{p}_k^2 + \hat{v}(\hat{R}_1) + \frac{m}{2} \sum_j \omega_j^2 (\hat{R}_j - \hat{R}_1)^2, \\ \hat{v}(\hat{R}_1) &\approx \frac{m}{2} \sum_{\pm} \Omega^2 (\hat{R}_1 \pm a)^2 \pm \frac{\Delta}{2}.\end{aligned}\tag{7}$$

The states for the two situations ”object in the left well” and ”object in the right well” will be denoted by $|\Lambda_L, L\rangle$ and $|\Lambda_R, R\rangle$, respectively. If one projects onto the eigenstates $|n\rangle$ of the one-dimensional harmonic oscillator and takes into account the shift of the wells, one arrives at the following decomposition ($\varphi_n(R) = \langle R|n\rangle$):

$$\langle n_L, \{R_j\} | \Lambda_L, L \rangle =: \int dR_1 \varphi_{n_L}(R_1 + a) \phi_{\Lambda_L}^L(\{R_k\}),\tag{8}$$

$$\langle n_R, \{R_j\} | \Lambda_R, R \rangle =: \int dR_1 \varphi_{n_R}(R_1 - a) \phi_{\Lambda_R}^R(\{R_k\}).\tag{9}$$

The situations ”object on the left” and ”object on the right” differ only by the shift and the bias of the wells. Therefore, one can find a unified representation by noting that $\phi_{\Lambda}^L(\{R_k\}) = \Phi_{\Lambda}(\{R_k + a\})$ and $\phi_{\Lambda}^R(\{R_k\}) = \Phi_{\Lambda}(\{R_k - a\})$. The eigenstates Φ_{Λ} are defined by the relations

$$\Phi_{\Lambda}(\{R_k\}) := \langle \{R_k\} | \Lambda \rangle,\tag{10}$$

$$\hat{H}_0 |\Lambda\rangle = E_{\Lambda} |\Lambda\rangle,\tag{11}$$

$$\hat{H}_0 = \frac{1}{2m} \sum_k \hat{p}_k^2 + \frac{m}{2} \sum_j \omega_j^2 (\hat{R}_j - \hat{R}_1)^2 + \frac{m}{2} \Omega^2 \hat{R}_1^2.\tag{12}$$

Thus, it follows from eqs. (8) and (9), respectively, that

$$\langle n_L, \{R_j\} | \Lambda_L, L \rangle = \langle n_L, \{R_j\} | \exp \left(ia \sum_j \hat{p}_j \right) | \Lambda_L \rangle, \quad (13)$$

$$\langle n_R, \{R_j\} | \Lambda_R, R \rangle = \langle n_R, \{R_j\} | \exp \left(-ia \sum_j \hat{p}_j \right) | \Lambda_R \rangle, \quad (14)$$

where we have used the shift property of the momentum operator $\hat{p}$.

The coupling of the two wells is taken into account by means of a tunneling Hamiltonian $\hat{H}_T$ which we represent in the form

$$\langle \Lambda_L, L | \hat{H}_T | \Lambda_R, R \rangle = \int d\{R_j\} \sum_{n_L n_R} T_{n_L n_R} \langle \Lambda_L, L | n_L, \{R_j\} \rangle \langle n_R, \{R_j\} | \Lambda_R, R \rangle. \quad (15)$$

Using again the momentum operator, one can write

$$|x\rangle \langle x'| = e^{i\hat{p}(x'-x)} |x'\rangle \langle x| = e^{i\hat{p}(x'-x)} \delta(x' - \hat{x}). \quad (16)$$

From this, we conclude that

$$\begin{aligned} \langle \Lambda_L, L | \hat{H}_T | \Lambda_R, R \rangle &= \sum_{n_L n_R} T_{n_L n_R} \int dR_1 dR'_1 \frac{dQ}{2\pi} \varphi_{n_R}^*(R'_1) \varphi_{n_L}(R_1) \\ &\times \langle \Lambda_L, L | e^{i\hat{p}_1(R'_1 - R_1)} e^{iQ(R'_1 - \hat{R}_1)} | \Lambda_R, R \rangle. \end{aligned} \quad (17)$$

III. TRANSITION RATE

The net transition rate from the left well to the right one is then in second order perturbation theory given by

$$\begin{aligned} J &= 2\pi Z_0^{-1} \sum_{\Lambda_L, \Lambda_R} |\langle \Lambda_L, L | \hat{H}_T | \Lambda_R, R \rangle|^2 \delta(E_{\Lambda_L} - E_{\Lambda_R} + \Delta) \\ &\times [e^{-\beta E_{\Lambda_L}} - e^{\beta E_{\Lambda_R}}], \end{aligned} \quad (18)$$

where $Z_0 = \text{Tr} \exp(-\beta H_0)$. The δ -function may be written in Fourier representation, and the fact that the E_Λ are eigenenergies of $\hat{H}_0$ serves us to incorporate the energy conservation into Heisenberg time-dependent operators $\hat{A}(t) = \exp(i\hat{H}_0 t) \hat{A} \exp(-i\hat{H}_0 t)$, i.e.

$$\begin{aligned}
& \langle \Lambda_L, L | \hat{H}_T | \Lambda_R, R \rangle \delta(E_{\Lambda_L} - E_{\Lambda_R} + \Delta) = \\
& = \int dt e^{i\Delta t} \langle \Lambda_L | e^{-ia \sum_k \hat{p}_k(t)} \hat{H}_T(t) e^{-ia \sum_k \hat{p}_k(t)} | \Lambda_R \rangle.
\end{aligned} \tag{19}$$

Then, collecting our results from eqs. (13), (14), (17), and (19), we arrive at the expression

$$\begin{aligned}
J &= Z_0^{-1} (1 - e^{-\beta\Delta}) \int dt e^{i\Delta t} \sum_{n_L, n_R} \sum_{\bar{n}_L, \bar{n}_R} T_{n_L, n_R} T_{\bar{n}_L, \bar{n}_R}^* \\
&\times \int \frac{dQ d\bar{Q}}{(2\pi)^2} \int dR_1 dR'_1 d\bar{R}_1 d\bar{R}'_1 \varphi_{n_L}(R_1) \varphi_{n_R}^*(R'_1) \varphi_{\bar{n}_L}^*(\bar{R}'_1) \varphi_{\bar{n}_R}(\bar{R}_1) \\
&\times \text{Tr} \{ e^{-\beta\hat{H}_0} e^{-2ia \sum_k \hat{p}_k(t)} e^{i\hat{p}_1(t)(R'_1 - R_1 + 2a)} e^{-iQ(\hat{R}_1(t) - R'_1)} \\
&\times e^{+2ia \sum_k \hat{p}_k} e^{i\hat{p}_1(\bar{R}'_1 - \bar{R}_1 - 2a)} e^{-i\bar{Q}(\hat{R}_1 - \bar{R}'_1)} \}.
\end{aligned} \tag{20}$$

Let us now use the relation

$$e^{-i(\hat{H}_0 + \hat{W})t} = e^{-i\hat{H}_0 t} \hat{T} e^{-i \int_0^t dt' \hat{W}(t')} \tag{21}$$

which holds for $t > 0$ when $\hat{T}$ is the time-ordering operator and for $t < 0$ when the anti time-ordering is used. If we define $\langle \hat{A} \rangle := \text{Tr} \exp(-\beta\hat{H}_0) \hat{A} / Z_0$, we can write the following result for the transition rate:

$$\begin{aligned}
J &= (1 - e^{-\beta\Delta}) \int dt e^{i(\Delta - 2m\Omega^2 a^2)t} \sum_{n_L, n_R} \sum_{\bar{n}_L, \bar{n}_R} T_{n_L, n_R} T_{\bar{n}_L, \bar{n}_R}^* \\
&\times \int \frac{dQ d\bar{Q}}{(2\pi)^2} \int dR dR' d\bar{R} d\bar{R}' e^{iQ \frac{R+R'}{2} + i\bar{Q} \frac{\bar{R}+\bar{R}'}{2}} \\
&\times \varphi_{n_L}(R) \varphi_{n_R}^*(R' - 2a) \varphi_{\bar{n}_L}^*(\bar{R}') \varphi_{\bar{n}_R}(\bar{R} - 2a) \\
&\times \left\langle \hat{T} \exp[-iQ\hat{R}_1(t) + i\hat{p}_1(t)(R' - R) + 2im\Omega^2 a \int_0^t dt' \hat{R}_1(t') \right. \\
&\quad \left. - i\bar{Q}\hat{R}_1(0) + i\hat{p}_1(0)(\bar{R}' - \bar{R})] \right\rangle.
\end{aligned} \tag{22}$$

We are now in the position to make use of the fact that the Hamiltonian is quadratic in all coordinates so that we can evaluate exactly

$$\langle \hat{T} e^{i \int dt' \eta(t') \hat{R}_1(t')} \rangle = e^{-\frac{i}{2} \int \int dt' dt'' \eta(t') D(t', t'') \eta(t'')}, \tag{23}$$

$$D(t', t'') := -i \langle \hat{T} \hat{R}_1(t') \hat{R}_1(t'') \rangle. \tag{24}$$

By comparison with the last two lines in eq. (22), the function $\eta(t')$ is given by

$$\begin{aligned}\eta(t') = & -Q\delta(t' - t) - \overline{Q}\delta(t') + 2m\Omega^2 a[\Theta(t') - \Theta(t' - t)] \\ & + m(R - R')\delta'(t' - t) + m(\overline{R} - \overline{R}')\delta'(t').\end{aligned}\quad (25)$$

$\Theta(t)$ is meant to represent the step function. The derivatives of the δ -function arise from a partial integration of terms containing $\hat{p}(t) = m d\hat{x}(t)/dt$. Note, that these act only on the coordinates but not on the step functions which arise due to the time ordering.

Moreover, the degrees of freedom of the bath can be integrated out in the usual way [2] leading to a dissipative influence on the object. One is then lead to the following form of the Fourier transform of $D(t, t') \equiv D(t - t')$:

$$D(\omega) = \frac{D^R(\omega)}{1 - \exp(-\hbar\omega/k_B T)} + \frac{D^R(-\omega)}{1 - \exp(\hbar\omega/k_B T)}, \quad (26)$$

$$(D^R)^{-1}(\omega) = m[(\omega + i0)^2 - \Omega^2] + i\eta\omega, \quad (27)$$

where we will use a spectral density $J(\omega) = \eta\omega$, $0 \leq \omega \leq \omega_c$ for the bath oscillators (see also [11] and appendix A).

From eqs. (22) and (23) one can conclude that the integrations with respect to $Q, \overline{Q}, R, R', \overline{R}, \overline{R}'$ can be done exactly as only Gaussian integrals are involved (note, that the eigenstates of the harmonic oscillator are Gaussian functions and derivatives of these, respectively). Therefore, for given $n_L, n_R, \overline{n}_L, \overline{n}_R$, one has to perform a 6-dimensional Gaussian integral. The only difficulty lies in the t -integration, which we have done numerically. Note furthermore, that without dissipation (i.e. $\eta = 0$) the tunneling current J is independent of the shift a , as one can show explicitly.

IV. TRANSITION RATE AT RESONANCE

Specifically, let us restrict ourselves to the case of zero temperature where only the lowest level of the left well is expected to contribute to the transition rate. Moreover, we will assume that the bias $\Delta \approx \Omega$ and that the dissipation leads only to a moderate broadening of the

energy levels of the oscillator. Then, one needs to consider only the levels $n_L = \bar{n}_L = 0$ and $n_R = \bar{n}_R = 1$.

The calculation of the transition rate for this case is deferred to appendix B. We want to mention, however, that the cutoff frequency ω_c of the bath spectrum is important to avoid a divergency of $\ddot{D}(t)$ for $t \rightarrow 0$. In fig. 2, we show the results for the transition rate calculated for different values of the dissipation η and the shift a . There, we have used dimensionless variables which means that the frequencies and lengths were normalized to their characteristic scales $\Omega = \sqrt{2eI_J/\hbar C}$ and $\kappa^{-1} = \sqrt{\hbar/M\Omega}$, respectively. The dimensionless parameters are then given by $a^* = \kappa a$ for the shift and $\eta^* = 1/(RC\Omega)$ for the viscosity.

In a typical experimental situation, one has $I_J \approx 1\text{nA}$ for the Josephson current and $C \approx 1\text{fF}$ for the capacity. Thus, we have $\kappa^{-1} \approx 4.2$, which means that $a^* \simeq \pi\kappa \approx 0.75$. The dimensionless viscosity is given by $\eta^* \approx 18\text{k}\Omega/R$. Note, however, that the transition rate to be observed experimentally will be very small due to the exponentially small factor $|T_{01}|^2/(\hbar\Omega)^2$ [8].

For small values of the dissipation or rather of $\alpha = 2\eta a^2/\pi$, the perturbation theory [10] yields the expression

$$J = 2|T_{01}|^2 \text{Re} \left(e^{i\pi\alpha} \left(\frac{\gamma + i\epsilon}{\omega_c} \right)^{2\alpha} \frac{1}{\gamma + i\epsilon} \right), \quad (28)$$

where $\gamma = \eta/(2m)$ and $\epsilon = \Delta - \Omega$. When normalized to the same value of the peak maximum, the curves calculated from the above formula (28) are nearly identical with those we obtained from our theory (cf. fig. 3). The different normalization stems from [12] the way the divergencies were removed in [10].

V. CONCLUSION

In summary, we have studied the transition rate in a double-well system under the influence of a dissipative environment. The rate shows asymmetric peaks when the levels of the wells approach each other. Thus, we have calculated the I-V characteristics of a

Josephson junction shunted by an Ohmic resistor for resonance conditions. We have modeled the system by two parabolic potentials coupled to a bath of harmonic oscillators. The coupling of the wells has been realized by a tunneling Hamiltonian. Moreover, the analysis has taken into account that the transfer of the object between the wells is a multiphonon process. In addition, our calculation has the advantage that the bath coordinates do not appear explicitly any more when the thermal average is taken.

ACKNOWLEDGMENTS

The authors would like to thank A.I. Larkin and Yu.N. Ovchinnikov for helpful discussions.

APPENDIX A: GREEN'S FUNCTIONS FOR THE HARMONIC OSCILLATOR IN A DISSIPATIVE ENVIRONMENT

From the equations of motion of the operators $\hat{R}_k(t)$ and $\hat{p}_k(t)$ in the Heisenberg representation, one can immediately infer the form of the retarded Green's function $D^R(t - t') = -i\Theta(t - t')\langle[\hat{x}(t), \hat{x}(t')]\rangle$:

$$(D^R)^{-1}(\omega) = m[(\omega + i0)^2 - \Omega^2] - \frac{2}{\pi} \int_0^\infty d\omega' \frac{\omega^2}{(\omega + i0)^2 - \omega'^2} \frac{J(\omega')}{\omega'} \quad (\text{A1})$$

$$= m[(\omega + i0)^2 - \Omega^2] + i\eta\omega \quad (\text{A2})$$

if the spectral density [3] is given by

$$J(\omega) = \frac{\pi}{2} \sum_j m\omega_j^3 \delta(\omega - \omega_j) = \eta\omega, \quad 0 \leq \omega \leq \omega_c. \quad (\text{A3})$$

In order to avoid divergencies of $\ddot{D}(t \rightarrow 0)$, it is necessary to introduce explicitly a cutoff for $J(\omega)$ with the characteristic frequency ω_c . We used a Lorentzian cutoff which permits to evaluate the integral in (A1) exactly. For $\eta/(2m) \ll \Omega \ll \omega_c$ we obtain

$$mD(t) = i \operatorname{Im} (Ae^{-i\tilde{\Omega}t}) - i \operatorname{Im} A e^{-t(\omega_c - \bar{\eta}/m)}$$

$$\begin{aligned}
& -\frac{1}{\pi} \text{Im} \left[A(g(-\tilde{\Omega}t) + g(\tilde{\Omega}t)) \right] \\
& + \frac{2}{\pi} \text{Im} A \text{Re} g(i\omega_c t - i\bar{\eta}t/m),
\end{aligned} \tag{A4}$$

where we have introduced

$$A = \frac{\tilde{\Omega} + i\omega_c}{2\bar{\Omega}(\tilde{\Omega} + i\omega_c - i\bar{\eta}/m)}, \tag{A5}$$

$$\tilde{\Omega} = \bar{\Omega} - i\bar{\eta}/(2m), \tag{A6}$$

$$\bar{\Omega} = \Omega \left(1 + \frac{\eta}{2m} \frac{\omega_c}{\omega_c^2 + \Omega^2} \right), \tag{A7}$$

$$\bar{\eta} = \eta \left(1 - \frac{\Omega^2}{\Omega^2 + \omega_c^2} \right). \tag{A8}$$

The function $g(\omega)$ is related to the exponential integral [13].

Finally, the correlation function $D^K(t - t') = -i\langle [\hat{x}(t), \hat{x}(t')]_+ \rangle$ is given by

$$D_\omega^K = 2i \coth \frac{\hbar\omega}{2k_B T} \text{Im} D_\omega^R, \tag{A9}$$

so that we have

$$2D(t) = D^R(t) + D^R(-t) + D^K(t), \tag{A10}$$

which immediately leads to eq. (26).

APPENDIX B: EVALUATION OF THE 6-DIMENSIONAL GAUSSIAN INTEGRAL

First, let us write the transition rate (22) in the form

$$J = 2 \text{Re} \int_0^\infty dt e^{i\Delta t} K(t). \tag{B1}$$

Also, we introduce new integration variables $\boldsymbol{x} = (v, q, u, v', q', u')$ such that

$$R = (u - v - u' + v')/4 + a,$$

$$R' = (u + v - u' - v')/4 + a,$$

$$\begin{aligned}
\overline{R} &= (u + v + u' + v')/4 + a, \\
\overline{R}' &= (u - v + u' - v')/4 + a, \\
Q &= (q - q')/2, \\
\overline{Q} &= (q + q')/2.
\end{aligned} \tag{B2}$$

In addition, the eigenstates of the harmonic oscillator are given by

$$\varphi_0(R) = \sqrt{\kappa\pi}^{-1/4} e^{-\kappa^2 R^2/2}, \tag{B3}$$

$$\varphi_1(R) = \sqrt{2}R\kappa \varphi_0(R), \quad \text{with} \tag{B4}$$

$$\kappa = \sqrt{m\Omega}. \tag{B5}$$

Thus, the expression for $K(t)$, $t \geq 0$, reads

$$\begin{aligned}
K(t) &= \int \frac{dq dq'}{4\pi^2} du du' dv dv' |T_{01}|^2 \frac{\kappa^2}{16\pi} \\
&\quad \times \frac{\partial^2}{\partial \overline{a} \partial \overline{a}'} \exp(-\mathbf{x} \mathbf{M} \mathbf{x} + \mathbf{b} \mathbf{x} + c) \Big|_{\overline{a}=0=\overline{a}'}.
\end{aligned} \tag{B6}$$

The only nonzero elements of the symmetric 6×6 matrix $\mathbf{M}$ are given by

$$\begin{aligned}
M_{11} &= \frac{\kappa^2}{4} - \frac{im^2}{2}(\ddot{D}(0) - \ddot{D}(t)), \\
M_{22} &= \frac{i}{2}(D(0) + D(t)), \\
M_{33} &= \frac{\kappa^2}{4} = M_{66}, \\
M_{12} &= M_{21} = -\frac{im}{2}\dot{D}(t) = -M_{45} = -M_{54}, \\
M_{23} &= M_{32} = -\frac{i}{4} = M_{56} = M_{65}, \\
M_{44} &= \frac{\kappa^2}{4} - \frac{im^2}{2}(\ddot{D}(0) + \ddot{D}(t)), \\
M_{55} &= \frac{i}{2}(D(0) - D(t)).
\end{aligned} \tag{B7}$$

Thus, it can be decomposed into two 3×3 blocks. Furthermore, the vector $\mathbf{b}$ reads

$$\begin{aligned}
\mathbf{b} &= \frac{i\kappa\overline{a}}{4}(1, 0, 1, 1, 0, 1) - \frac{i\kappa\overline{a}'}{4}(1, 0, 1, -1, 0, -1) \\
&\quad + \left(\kappa^2 a - 2im^2\Omega^2 a[D(t) - D(0)], \quad ia + 2im\Omega^2 a \int_0^t dt' D(t'), \quad 0, 0, 0, 0 \right),
\end{aligned} \tag{B8}$$

and

$$c = -2\kappa^2 a^2 - i\kappa a(\bar{a} - \bar{a}') - i2m^2\Omega^4 a^2 \int_0^t dt_1 dt_2 D(t_1 - t_2) - i2m\Omega^2 a^2 t. \quad (\text{B9})$$

The Gaussian integral in eq. (B6) can be evaluated exactly and eventually, we obtain

$$K(t) = W(t) e^{S(t)}, \quad (\text{B10})$$

$$W(t) = \frac{\kappa^2}{8} |T_{01}|^2 (\det \mathbf{M})^{-1/2} \times \left\{ \frac{\partial \mathbf{b}}{\partial \bar{a}} (\mathbf{M}^{-1}) \frac{\partial \mathbf{b}}{\partial \bar{a}'} - \left[\mathbf{b} (\mathbf{M}^{-1}) \frac{\partial \mathbf{b}}{\partial \bar{a}} + \frac{\partial c}{\partial \bar{a}} \right]^2 \right\}_{\bar{a}=0=\bar{a}'}, \quad (\text{B11})$$

$$S(t) = \left\{ \frac{1}{2} \mathbf{b} (\mathbf{M}^{-1}) \mathbf{b} + c \right\}_{\bar{a}=0=\bar{a}'}. \quad (\text{B12})$$

In the case of zero dissipation, this expression is indeed independent of the shift a , as it can be shown exactly.

REFERENCES

- [1] A.J. Leggett, in: *The Lesson of Quantum Theory*, Niels Bohr Centenary Symposium 1985; ed. by J. de Boer, E. Dal, and O. Ulfbeck, (North Holland, Amsterdam 1986)
- [2] A.O. Caldeira and A.J. Leggett, Phys. Rev. Lett. **46** (1981) 211; Ann. Phys. (New York) **149** (1983) 374
- [3] A.J. Leggett, S. Chakravarty, A.T. Dorsey, M.P.A. Fisher, A. Chang, and W. Zwerger, Rev. Mod. Phys. **59** (1987) 1
- [4] M. Sasseti and U. Weiss, Phys. Rev. Lett. **65** (1990) 2262
- [5] N.C. van der Vaart et al., Phys. Rev. Lett. **74** (1995) 4702
- [6] J.M. Martinis, M.H. Devoret, and J. Clarke, Phys. Rev. Lett. **55** (1985) 1543
- [7] A.I. Larkin and Yu. N. Ovchinnikov, Sov. Phys. JETP **91** (1986) 318
- [8] M.H. Devoret, J.M. Martinis, and J. Clarke, Phys. Rev. Lett. **55** (1985) 1908
- [9] A.I. Larkin, Yu. N. Ovchinnikov, and A. Schmid, Physica **B 152** (1988) 266
- [10] Yu.N. Ovchinnikov and A. Schmid, Phys. Rev. B **50** (1994) 6332
- [11] A. Schmid, Ann. Phys. **173** (1987) 103
- [12] Yu.N. Ovchinnikov, private communication (1995)
- [13] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions* (Dover, New York 1972), formula 5.2.7

FIGURES




FIG. 1. Double well potential with shift a and bias Δ




FIG. 2. Transition rate $J^*/|T_{01}^*|^2$ as a function of the bias Δ^* for $T = 0$ (dimensionless, $\omega_c^* = 60$)




FIG. 3. Comparison of $J^*(\Delta^*)/|T_{01}^*|^2$ (solid line) with perturbation theory (crosses) (dimensionless, $\eta^* = 0.01$, $a^* = 4$, $\omega_c^* = 80$, $T = 0$)